

Hydrostatic and Nonhydrostatic Nested Modeling of Straits in the Philippines Archipelago

Dr. Patrick C. Gallacher
Naval Research Laboratory, Ocean Sciences Branch
Stennis Space Center, MS 39529
phone: (228) 688-5315; fax: (228) 688-4149; e-mail: gallacher@nrlssc.navy.mil

Award Number "N00014-10-WX20784"

LONG-TERM GOALS

This study utilizes nested nonhydrostatic models embedded in hydrostatic models to simulate and predict the submesoscale dynamics of straits at high spatial and temporal resolutions. The goal of this work is to understand the submesoscale dynamics of straits and the impact of these dynamics on the throughflow in the straits. The Navy requires the ability to forecast features and circulations forced by these dynamics on scales that impact naval operations, i.e. kilometers to meters.

OBJECTIVES

The primary objective is to understand the submesoscale dynamics in straits using nested nonhydrostatic models embedded in hydrostatic models. Specifically we will work

- To understand the effects and interactions of the primary forcing components:
 - Tides, especially the spring-neap tidal cycle and remotely versus locally generated tides,
 - Large scale circulation, particularly the Pacific to Indian ocean throughflow and its seasonal variability,
 - Winds, especially the Southeast Asian monsoon cycle,
- To establish the resolution (dx and dz) and the aspect ratio (dx/dz) required to accurately simulate submesoscale physics and structures,
- To determine the importance of accurate and detailed representation of topography and forcing, especially at open boundaries,
- To understand the impact of rotation on the flow in straits, this is particularly important to nonhydrostatic physics,
- To explore the impact of data assimilation in a nonhydrostatic model, especially for sparse and irregular data,
- To compare model and field observations both for planning and for assessment.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Hydrostatic and Nonhydrostatic Nested Modeling of Straits in the Philippines Archipelago			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Ocean Sciences Branch, Stennis Space Center, MS, 39529			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

APPROACH

We use a system of multiply nested nonhydrostatic model (NRL-MIT) domains which utilize hydrostatic models (NCOM and/or HYCOM) to provide open boundary conditions for the coarsest NRL-MIT domain. The NRL-MIT domains consist of the nonhydrostatic version of the MITgcm model wrapped in a suite of scripts that provide initial/restart fields, open boundary values and handle output in a series of segmented, parallel integrations that maximize cpu usage and the ratio of system to wall clock time. The forcing consists of surface fluxes from the NOGAPS and COAMPS operational nowcast/forecast systems and open boundary conditions from the NCOM and/or HYCOM nowcast/forecast systems. HYCOM forecasts with resolutions of up to 4 km may be available in the region in the next year or two (Harley Hurlburt, personal communication). The basic bathymetry will be the NRL DBDB2 (2 minute) bathymetry which we hope will be enhanced and improved with several additional bathymetry databases obtained during the DRI.

WORK COMPLETED

Work has been severely hampered by the inconsistencies between the available numerical bathymetry databases and the bathymetry reported by the cruises. We continue to work to improve the bathymetry and to verify and analyze results in the rotated coordinate system.

Implemented, Tested and Published NEW Open Boundary Conditions for nesting a Nonhydrostatic model in a Hydrostatic model.

- Formulated a combined open boundary condition consisting of
 - A Transport Correction Scheme (TCS) for incoming fluxes to conserve volume
 - A Flow Relaxation Scheme (FRS) to match outgoing and incoming fluxes
- Developed and tested methods using hindcasts of the Mindoro straits (Gallacher *et al.* 2011)

Exploring Next Generation Visualization and Analysis for Very Large Datasets.

- Visualization and data management for very large datasets constitutes one of the next big challenges for HP Computing.
- We have conducted a very large experiment which generated 3.5 TB of data.
 - The domain for this experiment uses 2,000 processors and $3500 \times 2400 \times 40 = 336$ Million grid points, more than 3 times the number of grid points used by Global NCOM.
- Under the auspice of ONR 321US we have partnered with Makai Engineering to determine the feasibility of visualizing such a large dataset.
 - To that end we have downloaded the data onto an external disk provided by ONR and sent it to Makai engineering for preliminary processing and visualization
- Collaborating with Dr. Sean Ziegler of PETT, we are devising and testing new methods for data mining and visualization of large data sets.
 - Developing new nondimensional parameters to locate important areas or significant features in the data sets.

- Utilizing and improving scaling methods such as WKB to normalize widely varying data values.

RESULTS

Hindcasts of the 3D nonhydrostatic flow through the Mindoro straits at 500m resolution are forced with data from NRL-MTRY's RELO COAMPS. The model data are compared with mooring data in the Mindoro straits and the Panay Straits. The results are in good agreement in the Mindoro straits (Figure 3). However, in the Panay straits differences in model and in situ bathymetries caused significant discrepancies in the comparisons (Figure 4). Work is ongoing to improve the bathymetry and the open boundary conditions and to verify the rotated values.

IMPACT/APPLICATIONS

Tactical scale or submesoscale forecasting in domains of 100 to 200 km will require nonhydrostatic modeling systems with resolutions of 100s of meters or less to correctly predict the NLIWs, turbulent regions, fronts, boils and small scale eddies. This project studies the dynamics of NLIWs, their interactions and their impact on the tactical environment. This work furthers the basic understanding of NLIWs and lays the foundation for future nonhydrostatic forecast systems.

RELATED PROJECTS

This project is synergistic with the following projects:

Internal Waves in Straits (IWISE) ONR DRI,

Effects of Non-Acoustic Noise on Multi-Sensor USW Networks, NRL 6.2 core

—This work is supported by the HPC project

Nonhydrostatic Modeling of Nonlinear Internal Waves and Turbulence

PUBLICATIONS

Gallacher, P. C., D. A. Hebert, M. R. Schaferkotter, Nesting a Nonhydrostatic Model in a Hydrostatic Model: The Boundary Interface , *Ocean Modelling*, **40**, 2, 190-198, 2011.